

PARTRIDGE-SCHWENKE ROVIBRATIONAL ENERGY LEVELS FOR WATER: SPECTROSCOPIC ASSIGNMENT AND COMPARISON WITH EXPERIMENTAL ENERGY LEVELS

SERGEI A. TASHKUN, TATIANA A. PUTILOVA and SEMEN MIKHAILENKO, *Laboratory of Theoretical Spectroscopy, Institute of Atmospheric Optics, 1, Akademicheskii Av., 634055 Tomsk, Russia.*

Unambiguous rovibrational spectroscopic assignment ($\nu_1, \nu_2, \nu_3, J, K_a, K_c$) up to $J, K_a=30$ of PS energy levels for the (000), (010), (100), (020) and (001) vibrational states of nine isotopic species (H_2^{16}O , H_2^{18}O , H_2^{17}O , HD^{16}O , HD^{18}O , HD^{17}O , D_2^{16}O , D_2^{18}O , D_2^{17}O) of the water molecule will be presented. PS levels were calculated with the help of an accurate isotope dependent potential energy surface by Partridge and Schwenke^a and a large basis set. PS levels will be compared with experimental levels collected from the literature. Differences between PS and BT2^b calculated levels for the H_2^{16}O isotopic species will be also given and discussed.

^aH. Partridge and D.W. Schwenke, *J. Chem. Phys.* **106**(11), 4618-4639 (1997)

^bR.J. Barber J. Tennyson, G.J. Harris and R.N. Tolchenov, *Mon. Not. R. Astron.* **368**, 1087-1094 (2006)