

IR SPECTROSCOPY AT THE CANADIAN LIGHT SOURCE: THE ν_{11} FUNDAMENTAL AND $\nu_{16} + \nu_{18} - \nu_{18}$ HOT BAND OF TRANS-ACROLEIN

HONG-YU SHI, LI-HONG XU, R.M. LEES, *Centre for Laser, Atomic and Molecular Sciences (CLAMS), Department of Physics, University of New Brunswick, Saint John, NB, Canada E2L 4L5*; D.W. TOKARYK, *CLAMS, Department of Physics, University of New Brunswick, Fredericton, NB, Canada E3B 5A3*; A.R.W. McKELLAR, *Steacie Institute for Molecular Sciences, National Research Council, Ottawa, ON, Canada K1A 0R6*; D.R.T. APPADOO, *Canadian Light Source, 101 Perimeter Road, University of Saskatchewan, Saskatoon, SK, Canada S7N 0X4*.

The ν_{11} fundamental and the $\nu_{16} + \nu_{18} - \nu_{18}$ hot band have been identified in the high-resolution Fourier transform spectrum of trans-acrolein ($\text{CH}_2=\text{CH}-\text{CH}=\text{O}$) recorded in the 10- μm region on the Far-Infrared beamline of the Canadian Light Source synchrotron in Saskatoon. The two bands are centered at 912 cm^{-1} and 957 cm^{-1} , respectively, with the excited ν_{11} state corresponding to the A' in-plane CH_2 -rocking mode and the $\nu_{16} + \nu_{18}$ state to the combination of the A'' out-of-plane CH_2 rock with the low-frequency (158 cm^{-1}) A'' C-C torsional mode. This extends our previous work on the ν_{14} and ν_{16} fundamentals centered at 993 and 959 cm^{-1} , leaving only the ν_{15} mode now to be assigned at high resolution in the 10- μm spectral region for this important atmospheric pollutant. The ν_{11} band is type a/b and the $\nu_{16} + \nu_{18} - \nu_{18}$ hot band is c -type, so that assignments could be confirmed by lower-state combination differences. The assigned transitions have been fitted to a Watson asymmetric rotor Hamiltonian, and molecular parameters for both states will be reported.