

DOPPLER-FREE TWO-PHOTON EXCITATION SPECTROSCOPY OF BENZENE IN A MOLECULAR BEAM

YOSUKE SEMBA and SHUNJI KASAHARA, *Molecular Photoscience Research Center, Kobe University, Kobe 657-8501, Japan*; MASAOKI BABA, *Graduate School of Science, Kyoto University, Kyoto 606-8502, Japan*.

The fluorescence quantum yield at the vibrational energy over 3000 cm^{-1} in the $S_1\ ^1B_{2u}$ state of benzene is dramatically decreased. This phenomenon is called "channel three" and has been studied extensively. We have observed the ultrahigh-resolution spectrum of the $S_1 \leftarrow S_0\ 1_0^214_0^1$ band ($E_{\text{excess}} = 3412\text{ cm}^{-1}$) and found the Zeeman splitting which explained to be originating from the magnetic moment of the S_1 state induced by mixing with S_2 state by J - L coupling.^{a b} In this work, we established the apparatus for the Doppler-free two-photon excitation (DFTPE) spectroscopy with a molecular beam and measured the $S_1 \leftarrow S_0\ 1_0^114_0^1$ and $1_0^214_0^1$ band of benzene. When we measured the DFTPE spectrum in a gas cell, many rotational lines were observed but the pressure broadening was remained. On the other hand, when we measured the DFTPE spectrum in a molecular beam, the influence of collision was decreased. The typical linewidth was 6 MHz. The high-resolution and collision-free condition is useful to investigate the perturbing region with high sensitivity and accuracy.

^aD. Y. Baek, J. Chen, J. Wang, A. Doi, S. Kasahara, M. Baba, and H. Katô, *Bul. Chem. Soc. Jpn.*, **79**, 75 (2006).

^bH. Katô, M. Baba, and S. Kasahara, *Bul. Chem. Soc. Jpn.*, **80**, 456 (2007).