

JET-COOLED LASER SPECTROSCOPY OF A JAHN-TELLER AND PSEUDO JAHN-TELLER ACTIVE MOLECULE: THE NITRATE RADICAL

MING-WEI CHEN, *Laser Spectroscopy Facility, Department of Chemistry, The Ohio State University, 120 W. 18th Avenue, Columbus, Ohio 43210*; KANA TAKEMATSU, MITCHIO OKUMURA, *Arthur Amos Noyes Laboratory of Chemical Physics, California Institute of Technology, Pasadena, California 91125*; and TERRY A. MILLER, *Laser Spectroscopy Facility, Department of Chemistry, The Ohio State University, 120 W. 18th Avenue, Columbus, Ohio 43210*.

Well-known as an important intermediate in atmospheric chemistry, the nitrate radical (NO_3) has been extensively studied both experimentally and theoretically. The three energetically lowest electronic states ($\tilde{X}^2A'_2$, \tilde{A}^2E'' , and \tilde{B}^2E') are strongly coupled by vibronic interactions and hence it is a textbook molecule for understanding the coupling between nearby potential energy surfaces. Such coupling has been treated in considerable detail theoretically.^a However, corresponding experimental characterization of the interaction is much less detailed. The experimental results primarily consist of IR measurements of vibrational transitions in the ground state.^{bc} In addition, the electronically forbidden $\tilde{A}-\tilde{X}$ transition has been observed in ambient temperature CRDS studies.^d To understand both the Jahn-Teller and pseudo Jahn-Teller coupling in the molecule, further measurements are required with different selection rules and/or higher resolution to resolve the rotational structures of different transitions. In our group, a high-resolution (source $\Delta\nu \approx 100$ MHz in NIR region), jet-cooled CRDS system^e can be applied to rotationally resolve the electronically forbidden $\tilde{A}-\tilde{X}$ transition. Furthermore, our high-resolution LIF/SEP system (source $\Delta\nu \approx 100$ MHz) can provide the direct, rotationally resolved measurements of the $\tilde{B}-\tilde{X}$ and $\tilde{B}-\tilde{A}$ transitions by operating in the LIF and SEP modes respectively. Such data can provide unambiguous spectral assignments in the \tilde{X} , \tilde{A} and \tilde{B} states.

^aJ. F. Stanton, *J. Chem. Phys.*, **126**, 134309 (2007)

^bK. Kawaguchi, E. Hirota, T. Ishiwata, and I. Tanaka, *J. Chem. Phys.*, **93**, 951 (1990)

^cK. Kawaguchi, T. Ishiwata, E. Hirota, and I. Tanaka, *Chem. Phys.*, **231**, 193 (1998)

^dA. Deev, J. Sommar, and M. Okumura, *J. Chem. Phys.*, **122**, 224305 (2005)

^eS. Wu, P. Dupré, and T. A. Miller, *Phys. Chem. Chem. Phys.*, **8**, 1682, (2006)