

1-PHENYLPYRROLE A TICT MOLECULE?.^a

JESSICA A. THOMAS, JUSTIN W. YOUNG, ADAM J. FLEISHER, CASEY L. CLEMENTS, and DAVID W. PRATT, *Department of Chemistry, University of Pittsburgh, Pittsburgh, PA 15260*; LEONARDO ALVAREZ-VALTIERRA, *Division de Ciencias e Ingenierias, Campus Leon, Universidad de Guanajuato, Leon, Gto. 37150, Mexico*.

1-Phenylpyrrole has been identified by several authors as a molecule that displays twisted intramolecular charge-transfer (TICT) character in the condensed phase. However, Suzuka and coworkers^b, using single vibronic level dispersed fluorescence, observed no evidence of TICT activity in the gas phase. In this work, 1-phenylpyrrole is studied by the high resolution methods previously applied to DMABN^c to determine the extent of its motion along different possible TICT coordinates when it absorbs light.

^aWork supported by NSF CHE-0615755

^bK. Okuyama, Y. Numata, S. Odawara, and I. Suzuka, *J. Chem. Phys.*, **109**, 7185 (1998) and references therein.

^cA. E. Nikolaev, G. Myszkiewicz, G. Berden, L. Meerts, J. F. Pfanstiel, and D. W. Pratt, *J. Chem. Phys.*, **122**, 084309 (2005).