

## MOLECULES WITH A SIX-FOLD BARRIER: MICROWAVE SPECTRUM OF TOLUENE

VADIM V. ILYUSHIN, *Institute of Radio Astronomy of NASU, Chervonopraporna 4, 61002 Kharkov, Ukraine*; ZBIGNIEW KISIEL, LECH PSZCZÓLKOWSKI, *Institute of Physics, Polish Academy of Sciences, Al. Lotników 32/46, 02-668 Warszawa, Poland*; HEINRICH MÄDER, *Institut für Physikalische Chemie, Christian-Albrechts Universität zu Kiel, Olshausenstrasse 40, D-24098, Kiel, Germany*; JON T. HOUGEN, *Optical Technology Division, National Institute of Standards and Technology, Gaithersburg, MD 20899-8441*.

Recent progress in analysis and fitting of transitions in the toluene microwave spectrum characterized by  $J \leq 30$ ,  $K_a \leq 12$ , and the free rotor quantum number  $|m| \leq 3$  will be presented. The analysis was carried out using a new program which is specifically designed for six-fold barrier molecules, and allows the user to select almost any symmetry-allowed torsion-rotation term for inclusion in the fitting Hamiltonian. The program is based on the theoretical framework developed by Sørensen and Pedersen<sup>a</sup> in their application of the Longuet-Higgins permutation-inversion group  $G_{12}$  to the microwave spectrum of  $\text{CH}_3\text{NO}_2$ . In the analysis we have used the published and rather extensive unpublished data from previous studies as well as recent new measurements of the toluene microwave spectrum in the 8 – 18 GHz range. A rather successful fit of over 350 toluene microwave transitions in the abovementioned quantum number range has been achieved. The fit is close to the estimated experimental accuracy and includes more than thirty intertorsional  $m = +3 \leftrightarrow m = -3$  transitions.

---

<sup>a</sup>G. O. Sørensen, T. Pedersen, *Studies in Physical and Theoretical Chemistry*, Elsevier, **23**, 219–236 (1983).