

## CORRELATED AB INITIO STUDY OF THE GROUND ELECTRONIC STATE OF THE $\text{H}_2\text{-O}_2^-$ COMPLEX

WAFAA M. FAWZY, *Department of Chemistry, Murray State University, Murray, KY 42071.*

The  $\text{H}_2\text{-O}_2^-$  ( $X^2\Pi_g$ ) complex has been examined using the coupled-cluster theory at the CCSD(T)/aug-cc-pVDZ and CCSD(T)/aug-cc-pVTZ levels. Electronic structure calculations show that the global minimum energy structure corresponds to a planar bent geometry with a well depth of  $1550\text{ cm}^{-1}$ . For this geometry, the distance between centers of masses of moieties of the complex is  $2.57\text{ \AA}$ , angstroms and the angles between the internuclear axes of the superoxide radical and the hydrogen molecule with respect to the axis that connects their centers of masses are  $104^\circ$  and  $165^\circ$ , respectively. These results indicate that the hydrogen molecule and the superoxide radical are held together by a strong hydrogen bond within the complex. Results of the current work will be discussed and compared to results of our recent ab initio study<sup>a</sup> of the  $\text{H}_2\text{-O}_2$  ( $X^3\Sigma_g^-$ ) complex.

---

<sup>a</sup>Wafaa M. Fawzy, in preparation for publication