

MICROWAVE-MICROWAVE DOUBLE RESONANCE INVESTIGATION OF THE H₂-N₂O VAN DER WAALS COMPLEX

JEN NICOLE LANDRY, WOLFGANG JÄGER, *Department of Chemistry, University of Alberta, Edmonton, AB, T6G 2G2, Canada*; YOSHIHIRO SUMIYOSHI AND YASUKI ENDO, *Department of Basic Science, Graduate School of Arts and Sciences, The University of Tokyo, Komaba, Meguro-ku, Tokyo 153-8902, Japan*.

The $J=2-1$ rotational transitions of *ortho*D₂-¹⁵N₂O and *para*D₂-¹⁵N₂O were measured using a microwave-microwave double resonance technique. The experimental setup included a pulsed molecular beam Fourier Transform microwave spectrometer for signal detection and a microwave synthesizer with a K band standard gain horn (frequency range 18 - 26.5 GHz) to generate and transmit the pump radiation. The experimental arrangement was such that the pump radiation propagation was perpendicular to the supersonic molecular expansion. In these experiments, the pump radiation was tuned into resonance with the $J_{K_a K_c}=1_{01}-0_{00}$ or $1_{11}-0_{00}$ transition. The resulting population transfer from the rotational ground state facilitated the observation of the $J=2-1$ transition by significantly enhancing the signal intensity. In addition, a brief update on the solvation studies of nitrous oxide with *para*H₂ molecules will be given.