

LASER INDUCED FLUORESCENCE SPECTROSCOPY OF THE SiNSi RADICAL II: IDENTIFICATIONS OF THE \tilde{A}^2A_1 , \tilde{B}^2B_1 , AND $\tilde{D}^2\Sigma_g^+$ STATES

CHIHAYA MOTOYOSHI, YOSHIHIRO SUMIYOSHI, YASUKI ENDO, *Department of Basic Science, The University of Tokyo, Tokyo 153-8902, Japan*; MASARU FUKUSHIMA and TAKAHSHI ISHIWATA, *Faculty of Information Sciences, Hiroshima City University, Hiroshima 731-3194, Japan*.

Among the LIF spectra of SiNSi observed in the near UV region, another band system with fluorescence lifetimes of 30 ns was identified. From detailed analyses of the rotational structures of the observed vibronic bands, they were assigned to the $\tilde{D}^2\Sigma_g^+ - X^2\Pi_g$ band system of SiNSi, which is a symmetry forbidden transition borrowing intensity from vibronic mixings. Dispersed fluorescence spectra from SVL levels of the \tilde{D} state show distinct features around 13,000 cm^{-1} above the ground vibronic state, which were assigned to transitions to two electronic states \tilde{A}^2A_1 and \tilde{B}^2B_1 , correlated to a $^2\Pi_u$ state at the limit of linearity. LIF spectra were observed in the corresponding region from 13,000 to 14,000 cm^{-1} to confirm the assignment. Dispersed fluorescence spectra from the visible transitions gave vibrational term energies for high lying bending vibrational levels in the ground state, showing that they are bent-linear transitions.