

SUPERFLUID EFFECTS IN PARA-H₂ CLUSTERS PROBED BY CO₂ ROTATION-VIBRATION TRANSITIONS

HUI LI, ROBERT J. LE ROY, PIERRE-NICHOLAS ROY, *Department of Chemistry, University of Waterloo, Waterloo, ON N2L 3G1, Canada*; A.R.W. MCKELLAR, *Steacie Institute for Molecular Sciences, National Research Council of Canada, Ottawa, ON K1A 0R6, Canada*.

The prospect of directly observing superfluidity in para-H₂ is a tantalizing but elusive goal. Like ⁴He, para-H₂ is a light zero-spin boson. However, H₂-H₂ intermolecular interactions, though weak, are stronger than He-He interactions, and hydrogen is a solid below about 14 K. This makes detection of superfluidity in bulk hydrogen problematical, to say the least. But there are still possibilities for para-H₂ in the form of clusters or in nano-confined environments, and superfluid transition temperatures as high as ~ 6 K have been predicted.^a Spectroscopic observations of (para-H₂)_N-CO₂ clusters^b were at first very difficult to interpret for $N > 5$. However, with the help of path integral Monte Carlo simulations and an accurate new H₂-CO₂ intermolecular potential surface^c which explicitly incorporates dependence on the CO₂ ν_3 asymmetric stretch, it is now possible to achieve a remarkably consistent picture of (para-H₂)_N-CO₂ clusters in the size range $N = 1 \sim 20$. By combining the experimental spectroscopic measurements and theoretical simulations, we determine the size evolution of the superfluid response of the CO₂-doped para-H₂ clusters, which peaks for the “magic” number $N = 12$.

^aV.L. Ginzburg and A.A. Sobyenin, *JETP Lett.* **15**, 343 (1972).

^bA.R.W. McKellar, Paper WH04, 63rd OSU International Symposium on Molecular Spectroscopy, June 16-20, 2008.

^cH. Li, P.-N. Roy, and R.J. Le Roy, *J. Chem. Phys.*, submitted.