

VIBRATIONAL ANALYSIS AND VALENCE FORCE FIELD FOR NITROTOLUENES, DIMETHYLANILINES AND SOME SUBSTITUTED METHYLBENZENES

B. VENKATRAM REDDY, *Department of Physics, Kakatiya University, Warangal-506 009, A.P., India*
Email: bvreddy67@yahoo.com; JAI KISHAN OJHA, Department of Physics, Government Degree College, Mancherla-504 208, A.P., India; G. RAMANA RAO, Department of Physics, Varada Reddy College of Engineering, Ananthasagar, Warangal-506 371, A.P., India.

The Fourier transform infrared (FTIR) and Raman spectra of 2-amino-4-nitro-toluene; 2-amino-5-nitrotoluene; 2,4-dimethylaniline; 2,5-dimethylaniline; 2,6-dimethylaniline; 1,2,4-trimethylbenzene; 1,3,5-trimethylbenzene and pentamethyl-benzene have been recorded in the range $4000-400\text{ cm}^{-1}$ and $4000-30\text{ cm}^{-1}$, respectively. A normal coordinate analysis was carried out for both in-plane and out-of-plane vibrations of these molecules using an 81-parameter modified valence force field. The force constants were refined using 251 frequencies of eight molecules in the Overlay least-square technique. The reliability of force constants was tested by making zero-order calculations for both in-plane and out-of plane vibrations for five related molecules. The potential energy distribution (PED) and eigen vectors calculated in the process were used to make unambiguous vibrational assignment of all the fundamentals.