HIGH-RESOLUTION SPECTROSCOPY AND PRELIMINARY GLOBAL ANALYSIS OF C–H STRETCHING VIBRATIONS OF $C_2H_4$ IN THE 3000 AND 6000 $CM^{-1}$ REGIONS

M. A. LORONO GONZALEZ, *Department of Chemistry, Universidad de Oriente, Cumaná 6101, Estado Sucre, Venezuela*; V. BOUDON, M. LOËTE, *Laboratoire Interdisciplinaire Carnot de Bourgogne, UMR 5209 CNRS-Université de Bourgogne, 9. Av. A. Savary, BP 47870, F-21078 Dijon Cedex, France*; M. ROTGER, M.-T. BOURGEOIS, *Groupe de Spectrométrie Moléculaire et Atmosphérique, CNRS UMR 6089, Moulin de la Housse, BP 1039, Cases 16-17, F-51687 Reims Cedex 2, France*; K. DIDRICHE, M. HERMAN, *Laboratoire de Chimie quantique et Photophysique, CP160/09, Faculté des Sciences, Université Libre de Bruxelles, 50 ave. Roosevelt, B-1050, Brussels, Belgium*; V. A. KAPITANOV, Yu. N. PONOMAREV, A. A. SOLODOV, A. M. SOLODOV, T. M. PETROVA, *V.E. Zuev Institute of Atmospheric Optics SB RAS,1, Zuev Square, Tomsk, 634921, Russia*.

Ethylene is a naturally occurring compound in ambient air that affects atmospheric chemistry and global climate. The $C_2H_4$ spectrum is available in databases only for the 1000 and 3000 $cm^{-1}$ ranges. In this work[a], the ethylene absorption spectrum was measured in the 6030-6250 $cm^{-1}$ range with the use of a high resolution Bruker IFS 125HR Fourier-spectrometer and a two-channel opto-acoustic spectrometer with a diode laser. As a secondary standard of wavelengths, the methane absorption spectrum was used in both cases. A preliminary analysis was realized thanks to the tensorial formalism developed by the Dijon group that is implemented in the XTDS software package[b]. We considered the two combination bands $\nu_5 + \nu_9$ and $\nu_5 + \nu_{11}$ as an interacting dyad. Parameters for the $\nu_9/\nu_{11}$ dyad were fitted simultaneously from a re-analysis of previously recorded supersonic expansion jet FTIR data, while parameters for the $v_5 = 1$ Raman level were taken from literature. More than 600 lines could be assigned in the 6030-6250 $cm^{-1}$ region (and also 682 in the 2950–3150 $cm^{-1}$ region) and effective Hamiltonian parameters were fitted, including Coriolis interaction parameters. The dyad features are globally quite well reproduced, even if there are still problems at high $J$ values.

[a] M. A. Loroñno Gonzalez *et al.*, *J. Quant. Spectrosc. Radiat. Transfer*, **111**, 2265–2278 (2010).

[b] Ch. Wenger, V. Boudon, M. Rotger, J.-P. Champion and M. Sanzharov, *J. Mol. Spectrosc.*, **251**, 102–113 (2008).