

## UV AND IR SPECTROSCOPIC STUDIES OF COLD ALKALI METAL ION-BENZO CROWN ETHER COMPLEXES IN THE GAS PHASE

YOSHIYA INOKUCHI, *Department of Chemistry, Faculty of Science, Hiroshima University, Higashi-Hiroshima 739-8526, Japan*; OLEG V. BOYARKIN, *Laboratoire de Chimie Physique Moléculaire, École Polytechnique Fédérale de Lausanne, Lausanne CH-1015, Switzerland*; RYOJI KUSAKA, *Department of Chemistry, Faculty of Science, Hiroshima University, Higashi-Hiroshima 739-8526, Japan*; TAKEHARU HAINO, *Department of Chemistry, Faculty of Science, Hiroshima University, Higashi-Hiroshima 739-8526, Japan*; TAKAYUKI EBATA, *Department of Chemistry, Faculty of Science, Hiroshima University, Higashi-Hiroshima 739-8526, Japan*; and THOMAS R. RIZZO, *Laboratoire de Chimie Physique Moléculaire, École Polytechnique Fédérale de Lausanne, Lausanne CH-1015, Switzerland*.

We measured UV photodissociation and IR-UV double-resonance spectra of benzo-crown ether complexes with alkali metal ions ( $\text{Li}^+$ ,  $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Rb}^+$ , and  $\text{Cs}^+$ ) in a cold, 22-pole ion trap<sup>a</sup>. All the complexes show a number of vibronically resolved UV bands in the 36000-38000  $\text{cm}^{-1}$  region. We used the IR-UV spectra in the CH stretching region to distinguish peaks in the UV spectra that belong to different conformers. The use of density functional theory allows us to determine the conformation of the complexes.

---

<sup>a</sup>Y. Inokuchi, O. V. Boyarkin, R. Kusaka, T. Haino, T. Ebata and T. R. Rizzo *J. Am. Chem. Soc.* **133**(31), 12256 July 2011.