

RENNER-TELLER COUPLING IN H_2S^+ : PARTITIONING THE ROVIBRONIC AND SPINORBIT COUPLING HAMILTONIAN

G. DUXBURY, *Department of Physics, SUPA, John Anderson Building, University of Strathclyde, 107 Rottenrow, Glasgow G4 0NG, Scotland, UK*; Ch. JUNGEN, *LAC, 1 Laboratoire Aimé Cotton du CNRS, Université de Paris-Sud, 91405 Orsay, France*; A. ALIJAH, *GSMA, UMR CNRS 6089, Université de Reims Champagne-Ardenne, B.P. 1039, 51687 Reims Cedex 2, France*.

The stretch-bender reference-frame^a was developed to allow the separation of large amplitude bending motion and symmetric stretching. It has been used to calculate vibrational resonances, the effects of spin-orbit coupling, and of overall rotation. Its use allows the block factorisation of the Renner-Teller interaction matrix. We wish to show the utility of this approach when two different approaches by Dixon and Duxbury,^b and Jungen and Merer^c, are used to minimise the effects of the large amplitude bending upon the Renner-Teller interaction. It also allows the effects of large amplitude motion on the rotational structure to be calculated, including the switchover from bent to linear behaviour.

^aJ. Chem. Phys. 108, 2336 (1998) and J. Mol. Spectrosc. 211, 7 (2002)

^bMolec. Phys. 43, 255 (1981)

^cMolec. Phys. 40, 1 (1981)