

## PRODUCTION OF A BEAM OF HIGHLY VIBRATIONALLY EXCITED CO USING PERTURBATIONS

N. BARTELS, T. SCHÄFER, J. HÜHNERT, A. M. WODTKE, *Georg August Universität Göttingen, Institut für Physikalische Chemie, Tammanstrasse 6, 37075 Göttingen, Germany*; and R. W. FIELD, *Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, MA 02139, USA*.

For many experimentalists (especially those, who are not spectroscopists), molecular perturbations are a curse, as they make assignments and analysis of spectral data more difficult. Nevertheless, they can also be a boon!

In this talk we will show how a molecular beam of CO in high vibrational states ( $v = 17, 18$ ) can be prepared by an optical pumping scheme that we call PUMP-PUMP-PERTURB and DUMP (P<sup>3</sup>D). P<sup>3</sup>D exploits the loaning, via spin-orbit perturbations, of the large oscillator strength of the 4th positive system,  $A^1\Pi \leftarrow X^1\Sigma^+$ , to the triplet manifold.

This allows some nominally spin-forbidden transitions to be exploited in multistep optical pumping schemes. The ability to *state-selectively* prepare CO in high vibrational states opens up new opportunities for molecular beam scattering experiments.

