

## OBSERVATION OF BLUE-DETUNED PHOTOASSOCIATION TO THE $2(0_g^+)$ STATE OF $^{85}\text{Rb}_2$ VIA REMPI

M. A. BELLOS, R. CAROLLO, D. RAHMLow, J. BANERJEE, M. BERMUDEZ, E. E. EYLER, P. L. GOULD, and W. C. STWALLEY, *Department of Physics, University of Connecticut, Storrs, CT 06269.*

We report photoassociation of ultracold atoms to vibrational levels blue of the  $^{85}\text{Rb}_2$   $5s+5p_{1/2}$  asymptote, in the previously-unobserved  $2(0_g^+)$  Hund's case (c) state that corresponds to the  $2^1\Sigma_g^+$  state in Hund's case (a). These excited-state ultracold molecules decay to the  $a^3\Sigma_u^+$  state and are detected by pulsed REMPI through the  $2^3\Sigma_g^+$  state. We also see an order of magnitude enhancement in the  $v' = 111$ ,  $J' = 5$  ro-vibrational level of the  $2(0_g^+)$  state and present evidence for resonant coupling between this level and the  $v' = 155$  of the  $2(1_g)$  state, seen earlier in photoassociative trap loss<sup>a</sup>. Following photoassociation to the observed levels of the  $2(0_g^+)$  state, spontaneous decay populates vibrational levels approximately halfway up the  $a^3\Sigma_u^+$  potential well, including levels  $v'' = 18$  through  $v'' = 24$ . This pathway complements the blue-detuned photoassociation technique previously developed by this group<sup>b</sup>, which accesses the bottom of the  $a$  state potential. This work is supported by the NSF and AFOSR.

---

<sup>a</sup>R. A. Cline, J. D. Miller, and D. J. Heinzen, *Phys. Rev. Lett.* **73**, 632 (1994)

<sup>b</sup>M. A. Bellos, *et. al.*, *Phys. Chem. Chem. Phys.* **13**, 18880 (2011)