

HIGH-RESOLUTION PHOTOELECTRON SPECTROSCOPIC STUDY OF CYCLOPROPENE (c-C₃H₄)

KONSTANTINA VASILATO, JULIE M. MICHAUD, GUIDO GRASSI, DENITSA BAYKUSHEVA and
FREDERIC MERKT, *Laboratorium für Physikalische Chemie, ETH Zürich, CH-8093 Zürich, Switzerland.*

The spectroscopic information available on the cyclopropene radical cation is limited to that contained in low-resolution He I photoelectron spectra^{a,b}. To better characterize the structure of this cation, we have recently measured high-resolution PFI-ZEKE photoelectron spectra of c-C₃H₄ and several of its deuterated isotopomers in the vicinity of the adiabatic ionization threshold. Our new data include fully rotationally resolved spectra of the origin band of the $\tilde{X}^+ \leftarrow \tilde{X}$ transition of c-C₃H₄ and spectra of the low-vibrational levels of c-C₃H₄ and the deuterated isotopomers recorded at lower resolution. Because our efforts at synthesizing the partially deuterated isotopomers always resulted in mixtures of several isotopomers, differing in their number of D atoms and in the location of these atoms, the analysis of the isotopic shifts turned out to be challenging. Combining the information contained in the rotational structure of the origin band of c-C₃H₄ with the ionization energies of the isotopomers measured by photoionization mass spectrometry and the vibrational structure observed in the PFI-ZEKE spectra of the mixtures of deuterated isotopomers, we were able to draw conclusions on the structure of the cyclopropene radical cation based solely on experimental data. The adiabatic ionization energy of c-C₃H₄ was determined to be 77931.8(5) cm⁻¹.

^aM. B. ROBIN, C. R. BRUNDLE, N. A. KUEBLER, G. B. ELLISON AND K. B. WIBERG, *J. Chem. Phys.* **57**, 1758 (1972).

^bP. BISCHOF AND E. HEILBRONNER, *Helv. Chim. Acta* **53**, 1677 (1970).