

LINE BY LINE SPECTRAL PARAMETERS IN THE $4\nu_3$ SPECTRAL REGION OF METHANE

D. CHRIS BENNER, V. MALATHY DEVI, *Department of Physics, College of William and Mary, Williamsburg, VA 23187-8795*; J. J. O'BRIEN, S. SHAJI, *Department of Chemistry, University of Missouri - St. Louis, St. Louis, MO 63121-4400*; P. T. SPICKLER, C. P. HOUCK, J. A. COAKLEY, J. DOLPH, K. RANKIN, *Department of Physics, Bridgewater College, Bridgewater, VA 22812*.

The near infrared bands of methane were first observed in the outer planets and Titan where atmospheric ray paths are long. The spectrum is complex, and long absorption paths in the laboratory are difficult to cool to outer solar system temperatures. At room temperature, many significant spectral lines appear per Doppler width. The band models generally used in the 890 nm spectral region of methane do not provide transmissions that are multiplicative, so scattering and inhomogeneous atmospheres cannot be properly treated using this approach.

The intracavity laser spectrometer at the University of Missouri-St. Louis was used to obtain low temperature (99-161K), low pressure (0.12-7.13 Torr), long path (3.14-5.65 km) and high resolution (0.01 cm^{-1} HWHM) spectra of methane covering the entire 890nm feature (10925-11500 cm^{-1}), the deepest band in the CCD spectral region. At these temperatures the Doppler width is 0.01 cm^{-1} and the spectral lines originating from levels higher than $J''=11$ and excited vibrational states are not visible. The result is a dense, but manageable spectrum from which over 11,200 line positions, intensities and lower state energies are derived on a line by line basis by the College of William and Mary multispectrum nonlinear least squares fitting program^a. Simulations of the methane spectrum for outer planet atmospheres using our positions, intensities and lower state energies reveal a surprising amount of spectral structure at high resolution. This structure carries a great deal of atmospheric information^b.

^aD. Chris Benner, C. P. Rinsland, V. M. Devi, M. A. H. Smith, and D. A. Atkins, JQSRT 1995;53:705-21.

^bSupport for the work at William and Mary was provided by NASA through grant NNX08AF06G. Support for the work at UM-St. Louis provided by NASA through grant NAG5-12013, from NSF through grant CHE-0213356 and by the University of Missouri Research Board. Partial support at Bridgewater College was provided by its Martin Science Research Institute and from an AAS Small Research Grant.